


References

- [1] Torre LA, Bray F, Siegel RL et al. Global cancer statistics, 2012. *CA Cancer J Clin* 2015; 65: 87–108
- [2] van der Stok EP, Spaander MCW, Grünhagen DJ et al. Surveillance after curative treatment for colorectal cancer. *Nat Rev Clin Oncol* 2016; 14: 297
- [3] Levin TR, Corley DA, Jensen CD et al. Effects of organized colorectal cancer screening on cancer incidence and mortality in a large, community-based population. *Gastroenterology* 2018; 155: 1383–1391. e5
- [4] Logan RF, Patnick J, Nickerson C et al. Outcomes of the bowel cancer screening programme (BCSP) in England after the first 1 million tests. *Gut* 2012; 61: 1439–1446
- [5] Williams JG, Pullan RD, Hill J et al. Management of the malignant colorectal polyp: ACPGIBI position statement. *Colorectal Dis* 2013; 15 (Suppl. 02): 1–38
- [6] Richardson WS, Wilson MC, Nishikawa J et al. The well-built clinical question: A key to evidence-based decisions. *ACP J Club* 1995; 123: A12–A13
- [7] Atkins D, Best D, Briss PA et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004; 328: 1490
- [8] Spada C, Stoker J, Alarcon O et al. Clinical indications for computed tomographic colonography: European Society of Gastrointestinal Endoscopy (ESGE) and European Society of Gastrointestinal and Abdominal Radiology (ESGAR) Guideline. *Endoscopy* 2014; 46: 897–915
- [9] Kaminski MF, Thomas-Gibson S, Bugajski M et al. Performance measures for lower gastrointestinal endoscopy: a European Society of Gastrointestinal Endoscopy (ESGE) quality improvement initiative. *United European Gastroenterol J* 2017; 5: 309–334
- [10] Dumonceau JM, Hassan C, Riphaut A et al. European Society of Gastrointestinal Endoscopy (ESGE) Guideline Development Policy. *Endoscopy* 2012; 44: 626–629
- [11] Rosati G, Ambrosini G, Barni S et al. A randomized trial of intensive versus minimal surveillance of patients with resected Dukes B2-C colorectal carcinoma. *Ann Oncol* 2016; 27: 274–280
- [12] Rodriguez-Moranta F, Salo J, Arcusa A et al. Postoperative surveillance in patients with colorectal cancer who have undergone curative resection: a prospective, multicenter, randomized, controlled trial. *J Clin Oncol* 2006; 24: 386–393
- [13] Schoemaker D, Black R, Giles L et al. Yearly colonoscopy, liver CT, and chest radiography do not influence 5-year survival of colorectal cancer patients. *Gastroenterology* 1998; 114: 7–14
- [14] Makela JT, Laitinen SO, Kairaluoma MI. Five-year follow-up after radical surgery for colorectal cancer. Results of a prospective randomized trial. *Arch Surg* 1995; 130: 1062–1067
- [15] Secco GB, Fardelli R, Gianquinto D et al. Efficacy and cost of risk-adapted follow-up in patients after colorectal cancer surgery: a prospective, randomized and controlled trial. *Eur J Surg Oncol* 2002; 28: 418–423
- [16] Wattoo DA, Weller DP, Esterman A et al. General practice vs surgical-based follow-up for patients with colon cancer: randomised controlled trial. *Br J Cancer* 2006; 94: 1116–1121
- [17] Grossmann EM, Johnson FE, Virgo KS et al. Follow-up of colorectal cancer patients after resection with curative intent—the GILDA trial. *Surg Oncol* 2004; 13: 119–124
- [18] Pietra N, Sarli L, Costi R et al. Role of follow-up in management of local recurrences of colorectal cancer: a prospective, randomized study. *Dis Colon Rectum* 1998; 41: 1127–1133
- [19] Mokhles S, Macbeth F, Farewell V et al. Meta-analysis of colorectal cancer follow-up after potentially curative resection. *Br J Surg* 2016; 103: 1259–1268
- [20] Jeffery M, Hickey BE, Hider PN et al. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database Syst Rev* 2016; 11: Cd002200
- [21] Pita-Fernandez S, Alhayek-Ai M, Gonzalez-Martin C et al. Intensive follow-up strategies improve outcomes in nonmetastatic colorectal cancer patients after curative surgery: a systematic review and meta-analysis. *Ann Oncol* 2015; 26: 644–656
- [22] Tjandra JJ, Chan MK. Follow-up after curative resection of colorectal cancer: a meta-analysis. *Dis Colon Rectum* 2007; 50: 1783–1799
- [23] Wang T, Cui Y, Huang WS et al. The role of postoperative colonoscopic surveillance after radical surgery for colorectal cancer: a prospective, randomized clinical study. *Gastrointest Endosc* 2009; 69: 609–615
- [24] Choe EK, Park KJ, Chung SJ et al. Colonoscopic surveillance after colorectal cancer resection: who needs more intensive follow-up? *Digestion* 2015; 91: 142–149
- [25] Platell C, Salama P, Barwood N et al. Performing a colonoscopy 12 months after surgery for colorectal neoplasia. *ANZ J Surg* 2005; 75: 282–285
- [26] Barrier A, Houry S, Huguier M. The appropriate use of colonoscopy in the curative management of colorectal cancer. *Int J Colorectal Dis* 1998; 13: 93–98
- [27] Barillari P, Ramacciato G, Manetti G et al. Surveillance of colorectal cancer: effectiveness of early detection of intraluminal recurrences on prognosis and survival of patients treated for cure. *Dis Colon Rectum* 1996; 39: 388–393
- [28] Lautenbach E, Forde KA, Neugut AI. Benefits of colonoscopic surveillance after curative resection of colorectal cancer. *Ann Surg* 1994; 220: 206–211
- [29] Moon CM, Cheon JH, Choi EH et al. Advanced synchronous adenoma but not simple adenoma predicts the future development of metachronous neoplasia in patients with resected colorectal cancer. *J Clin Gastroenterol* 2010; 44: 495–501
- [30] Gresham G, Renouf DJ, Chan M et al. Association between palliative resection of the primary tumor and overall survival in a population-based cohort of metastatic colorectal cancer patients. *Ann Surg Oncol* 2014; 21: 3917–3923
- [31] 't Lam-Boer J, Van der Geest LG, Verhoef C et al. Palliative resection of the primary tumor is associated with improved overall survival in incurable stage IV colorectal cancer: A nationwide population-based propensity-score adjusted study in the Netherlands. *Int J Cancer* 2016; 139: 2082–2094
- [32] Mulder SA, Kranse R, Damhuis RA et al. The incidence and risk factors of metachronous colorectal cancer: an indication for follow-up. *Dis Colon Rectum* 2012; 55: 522–531
- [33] Green RJ, Metlay JP, Propert K et al. Surveillance for second primary colorectal cancer after adjuvant chemotherapy: an analysis of Inter-group 0089. *Ann Intern Med* 2002; 136: 261–269
- [34] le Clercq CM, Winkens B, Bakker CM et al. Metachronous colorectal cancers result from missed lesions and non-compliance with surveillance. *Gastrointest Endosc* 2015; 82: 325–333. e322
- [35] Liu L, Lemmens VE, De Hingh IH et al. Second primary cancers in subsites of colon and rectum in patients with previous colorectal cancer. *Dis Colon Rectum* 2013; 56: 158–168
- [36] Chen TA, Horng JT, Lin WC. Metachronous colorectal cancer in Taiwan: analyzing 20 years of data from Taiwan Cancer Registry. *Int J Clin Oncol* 2013; 18: 267–272
- [37] Couch DG, Bullen N, Ward-Booth SE et al. What interval between colorectal cancer resection and first surveillance colonoscopy? An audit of practice and yield. *Colorectal Dis* 2013; 15: 317–322
- [38] Chung YW, Han DS, Park KH et al. Patient factors predictive of inadequate bowel preparation using polyethylene glycol: a prospective study in Korea. *J Clin Gastroenterol* 2009; 43: 448–452

- [39] Hassan C, Fuccio L, Bruno M et al. A predictive model identifies patients most likely to have inadequate bowel preparation for colonoscopy. *Clin Gastroenterol Hepatol* 2012; 10: 501–506
- [40] Lim SW, Seo YW, Sinn DH et al. Impact of previous gastric or colonic resection on polyethylene glycol bowel preparation for colonoscopy. *Surg Endosc* 2012; 26: 1554–1559
- [41] Pontone S, Leonetti G, Lamazza A et al. A retrospective case-control study evaluating the bowel preparation quality during surveillance colonoscopy after colonic resection. *ISRN Gastroenterol* 2014; 2014: 681978
- [42] Pinol V, Andreu M, Castells A et al. Synchronous colorectal neoplasms in patients with colorectal cancer: predisposing individual and familial factors. *Dis Colon Rectum* 2004; 47: 1192–1200
- [43] Mulder SA, Kranse R, Damhuis RA et al. Prevalence and prognosis of synchronous colorectal cancer: a Dutch population-based study. *Cancer Epidemiol* 2011; 35: 442–447
- [44] Kaminski MF, Regula J, Kraszewska E et al. Quality indicators for colonoscopy and the risk of interval cancer. *NEJM* 2010; 362: 1795–1803
- [45] Rex DK, Bond JH, Winawer S et al. Quality in the technical performance of colonoscopy and the continuous quality improvement process for colonoscopy: recommendations of the U.S. Multi-Society Task Force on Colorectal Cancer. *Am J Gastroenterol* 2002; 97: 1296–1308
- [46] Rex DK, Petrini JL, Baron TH et al. Quality indicators for colonoscopy. *Am J Gastroenterol* 2006; 101: 873–885
- [47] Barclay RL, Vicari JJ, Doughty AS et al. Colonoscopic withdrawal times and adenoma detection during screening colonoscopy. *NEJM* 2006; 355: 2533–2541
- [48] Rex DK. Colonoscopic withdrawal technique is associated with adenoma miss rates. *Gastrointest Endosc* 2000; 51: 33–36
- [49] Cooper GS, Xu F, Barnholtz Sloan JS et al. Prevalence and predictors of interval colorectal cancers in medicare beneficiaries. *Cancer* 2012; 118: 3044–3052
- [50] Kaminski MF, Wieszczy P, Rupinski M et al. Increased rate of adenoma detection associates with reduced risk of colorectal cancer and death. *Gastroenterology* 2017; 153: 98–105
- [51] Lee RH, Tang RS, Muthusamy VR et al. Quality of colonoscopy withdrawal technique and variability in adenoma detection rates (with videos). *Gastrointest Endosc* 2011; 74: 128–134
- [52] Shaukat A, Rector TS, Church TR et al. Longer withdrawal time is associated with a reduced incidence of interval cancer after screening colonoscopy. *Gastroenterology* 2015; 149: 952–957
- [53] Atkin W, Wooldrage K, Brenner A et al. Adenoma surveillance and colorectal cancer incidence: a retrospective, multicentre, cohort study. *Lancet Oncol* 2017; 18: 823–834
- [54] Rutter MD, Senore C, Bisschops R et al. The European Society of Gastrointestinal Endoscopy Quality Improvement Initiative: developing performance measures. *United European Gastroenterol J* 2016; 4: 30–41
- [55] Mussetto A, Frazzoni L, Paggi S et al. Split dosing with a low-volume preparation is not inferior to split dosing with a high-volume preparation for bowel cleansing in patients with a history of colorectal resection: a randomized trial. *Endoscopy* 2015; 47: 917–924
- [56] Carballal S, Maisterra S, Lopez-Serrano A et al. Real-life chromoendoscopy for neoplasia detection and characterisation in long-standing IBD. *Gut* 2018; 67: 70–77
- [57] Rahmi G, Lecomte T, Malka D et al. Impact of chromoscopy on adenoma detection in patients with Lynch syndrome: a prospective, multicenter, blinded, tandem colonoscopy study. *Am J Gastroenterol* 2015; 110: 288–298
- [58] Bouvier AM, Latournerie M, Jooste V et al. The lifelong risk of metachronous colorectal cancer justifies long-term colonoscopic follow-up. *Eur J Cancer* 2008; 44: 522–527
- [59] Loberg M, Kalager M, Holme O et al. Long-term colorectal-cancer mortality after adenoma removal. *NEJM* 2014; 371: 799–807
- [60] Lee SY, Kim BC, Han KS et al. Incidence and risk factors of metachronous colorectal neoplasm after curative resection of colorectal cancer in Korean patients. *J Dig Dis* 2014; 15: 367–376
- [61] Patel A, Williams N, Parsons N et al. Risk factors for metachronous adenoma in the residual colon of patients undergoing curative surgery for colorectal cancer. *Int J Colorectal Dis* 2017; 32: 1609–1616
- [62] Fuccio L, Rex D, Ponchon T et al. New and recurrent colorectal cancers after resection: a systematic review and meta-analysis of endoscopic surveillance studies. *Gastroenterology* 2018; doi:10.1053/j.gastro.2018.12.006
- [63] Hyman N, Moore J, Cataldo P et al. The high yield of 1-year colonoscopy after resection: is it the handoff? *Surg Endosc* 2010; 24: 648–652
- [64] Rulyak SJ, Lieberman DA, Wagner EH et al. Outcome of follow-up colon examination among a population-based cohort of colorectal cancer patients. *Clin Gastroenterol Hepatol* 2007; 5: 470–476; quiz 407
- [65] Lan YT, Lin JK, Li AF et al. Metachronous colorectal cancer: necessity of post-operative colonoscopic surveillance. *Int J Colorectal Dis* 2005; 20: 121–125
- [66] Battersby NJ, Coupland A, Bouliotis G et al. Metachronous colorectal cancer: a competing risks analysis with consideration for a stratified approach to surveillance colonoscopy. *J Surg Oncol* 2014; 109: 445–450
- [67] Cone MM, Beck DE, Hicks TE et al. Timing of colonoscopy after resection for colorectal cancer: are we looking too soon? *Dis Colon Rectum* 2013; 56: 1233–1236
- [68] Fuccio L, Spada C, Frazzoni L et al. Higher adenoma recurrence rate after left- versus right-sided colectomy for colon cancer. *Gastrointest Endosc* 2015; 82: 337–343
- [69] McFall MR, Woods WG, Miles WF. Colonoscopic surveillance after curative colorectal resection: results of an empirical surveillance programme. *Colorectal Dis* 2003; 5: 233–240
- [70] Rajaratnam SG, Dennett ER. Development of metachronous neoplasms after colorectal cancer resection: absence of synchronous neoplasms predicts a lower risk. *N Z Med J* 2009; 122: 61–66
- [71] Sakamoto T, Matsuda T, Nakajima T et al. How often should we perform surveillance colonoscopy after surgery for colorectal cancer? *Int J Colorectal Dis* 2013; 28: 835–840
- [72] Jayasekara H, Reece JC, Buchanan DD et al. Risk factors for metachronous colorectal cancer following a primary colorectal cancer: A prospective cohort study. *Int J Cancer* 2016; 139: 1081–1090
- [73] Gervaz P, Bucher P, Neyroud-Caspar I et al. Proximal location of colon cancer is a risk factor for development of metachronous colorectal cancer: a population-based study. *Dis Colon Rectum* 2005; 48: 227–232
- [74] Marques-Antunes J, Libanio D, Goncalves P et al. Incidence and predictors of adenoma after surgery for colorectal cancer. *Eur J Gastroenterol Hepatol* 2017; 29: 932–938
- [75] Borda A, Martinez-Penuela JM, Borda F et al. Drawing up an individual risk index for development of metachronous neoplastic lesions in resected colorectal cancer. *Rev Esp Enferm Dig* 2012; 104: 291–297
- [76] Hollington P, Tiong L, Young G. Timing and detection of metachronous colorectal cancer. *ANZ J Surg* 2011; 81: 272–274
- [77] Kjeldsen B, Kronborg O, Fenger C et al. A prospective randomized study of follow-up after radical surgery for colorectal cancer. *Br J Surg* 1997; 84: 666–669
- [78] Park IJ, Yu CS, Kim HC et al. Metachronous colorectal cancer. *Colorectal Dis* 2006; 8: 323–327
- [79] Chen HS, Sheen-Chen SM. Synchronous and “early” metachronous colorectal adenocarcinoma: analysis of prognosis and current trends. *Dis Colon Rectum* 2000; 43: 1093–1099

- [80] Chen F, Stuart M. Colonoscopic follow-up of colorectal carcinoma. *Dis Colon Rectum* 1994; 37: 568–572
- [81] Granqvist S, Karlsson T. Postoperative follow-up of patients with colorectal carcinoma by colonoscopy. *Eur J Surg* 1992; 158: 307–312
- [82] Juhl G, Larson GM, Mullins R et al. Six-year results of annual colonoscopy after resection of colorectal cancer. *World J Surg* 1990; 14: 255–260; discussion 260–251
- [83] Togashi K, Konishi F, Ozawa A et al. Predictive factors for detecting colorectal carcinomas in surveillance colonoscopy after colorectal cancer surgery. *Dis Colon Rectum* 2000; 43: S47–S53
- [84] Kahi CJ, Boland CR, Dominitz JA et al. Colonoscopy surveillance after colorectal cancer resection: recommendations of the US Multi-Society Task Force on Colorectal Cancer. *Am J Gastroenterol* 2016; 111: 337–346; quiz 347
- [85] Fajobi O, Yiu CY, Sen-Gupta SB et al. Metachronous colorectal cancers. *Br J Surg* 1998; 85: 897–901
- [86] Papadopoulos V, Michalopoulos A, Basdanis G et al. Synchronous and metachronous colorectal carcinoma. *Tech Coloproctol* 2004; 8: (Suppl. 01): s97–s100
- [87] Hassan C, Pickhardt PJ, Zullo A et al. Cost-effectiveness of early colonoscopy surveillance after cancer resection. *Dig Liver Dis* 2009; 41: 881–885
- [88] Hojo K. Anastomotic recurrence after sphincter-saving resection for rectal cancer. Length of distal clearance of the bowel. *Dis Colon Rectum* 1986; 29: 11–14
- [89] Jung WB, Yu CS, Lim SB et al. Anastomotic Recurrence After Curative Resection for Colorectal Cancer. *World J Surg* 2017; 41: 285–294
- [90] Matsuda K, Hotta T, Takifuji K et al. Clinicopathological features of anastomotic recurrence after an anterior resection for rectal cancer. *Langenbecks Arch Surg* 2010; 395: 235–239
- [91] Obrand DI, Gordon PH. Incidence and patterns of recurrence following curative resection for colorectal carcinoma. *Dis Colon Rectum* 1997; 40: 15–24
- [92] de Chaisemartin C, Penna C, Goere D et al. Presentation and prognosis of local recurrence after total mesorectal excision. *Colorectal Dis* 2009; 11: 60–66
- [93] McDermott FT, Hughes ES, Pihl E et al. Local recurrence after potentially curative resection for rectal cancer in a series of 1008 patients. *Br J Surg* 1985; 72: 34–37
- [94] Patel SA, Chen YH, Hornick JL et al. Early-stage rectal cancer: clinical and pathologic prognostic markers of time to local recurrence and overall survival after resection. *Dis Colon Rectum* 2014; 57: 449–459
- [95] Kim Y-W, Kim N-K, Min B-S et al. Factors associated with anastomotic recurrence after total mesorectal excision in rectal cancer patients. *J Surg Oncol* 2009; 99: 58–64
- [96] Kang DW, Kwak HD, Sung NS et al. Oncologic outcomes in rectal cancer patients with a ≤ 1 -cm distal resection margin. *Int J Colorectal Dis* 2017; 32: 325–332
- [97] Stulc JP, Petrelli NJ, Herrera L et al. Anastomotic recurrence of adenocarcinoma of the colon. *Arch Surg* 1986; 121: 1077–1080
- [98] Pihl E, Hughes ES, McDermott FT et al. Recurrence of carcinoma of the colon and rectum at the anastomotic suture line. *Surg Gynecol Obstet* 1981; 153: 495–496
- [99] Harris GJ, Church JM, Senagore AJ et al. Factors affecting local recurrence of colonic adenocarcinoma. *Dis Colon Rectum* 2002; 45: 1029–1034
- [100] Bowne WB, Lee B, Wong WD et al. Operative salvage for locoregional recurrent colon cancer after curative resection: an analysis of 100 cases. *Dis Colon Rectum* 2005; 48: 897–909
- [101] Nash GM, Weiss A, Dasgupta R et al. Close distal margin and rectal cancer recurrence after sphincter-preserving rectal resection. *Dis Colon Rectum* 2010; 53: 1365–1373
- [102] Hogan J, Chang KH, Duff G et al. Lymphovascular invasion: a comprehensive appraisal in colon and rectal adenocarcinoma. *Dis Colon Rectum* 2015; 58: 547–555
- [103] Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? *J Clin Oncol* 2008; 26: 303–312
- [104] Adam IJ, Mohamdee MO, Martin IG et al. Role of circumferential margin involvement in the local recurrence of rectal cancer. *Lancet* 1994; 344: 707–711
- [105] Matsuda A, Kishi T, Musso G et al. The effect of intraoperative rectal washout on local recurrence after rectal cancer surgery: a meta-analysis. *Ann Surg Oncol* 2013; 20: 856–863
- [106] Zhou C, Ren Y, Wang K et al. Intra-operative rectal washout with saline solution can effectively prevent anastomotic recurrence: a meta-analysis. *Asian Pac J Cancer Prev* 2013; 14: 7155–7159
- [107] Terzi C, Unek T, Sagol O et al. Is rectal washout necessary in anterior resection for rectal cancer? A prospective clinical study *World J Surg* 2006; 30: 233–241
- [108] Docherty JG, McGregor JR, Akyol AM et al. Comparison of manually constructed and stapled anastomoses in colorectal surgery. West of Scotland and Highland Anastomosis Study Group. *Ann Surg* 1995; 221: 176–184
- [109] van den Brink M, Stiggelbout AM, van den Hout WB et al. Clinical nature and prognosis of locally recurrent rectal cancer after total mesorectal excision with or without preoperative radiotherapy. *J Clin Oncol* 2004; 22: 3958–3964
- [110] Ohlsson B, Breland U, Ekberg H et al. Follow-up after curative surgery for colorectal carcinoma. Randomized comparison with no follow-up. *Dis Colon Rectum* 1995; 38: 619–626
- [111] Figueredo A, Rumble RB, Maroun J et al. Follow-up of patients with curatively resected colorectal cancer: a practice guideline. *BMC Cancer* 2003; 3: 26
- [112] Renehan AG, Egger M, Saunders MP et al. Mechanisms of improved survival from intensive followup in colorectal cancer: a hypothesis. *Br J Cancer* 2005; 92: 430–433
- [113] Snyder RA, Hu C, Cuddy A et al. Association between intensity of posttreatment surveillance testing and detection of recurrence in patients with colorectal cancer. *JAMA* 2018; 319: 2104–2115
- [114] Wille-Jørgensen P, Syk I, Smedh K et al. Effect of more vs less frequent follow-up testing on overall and colorectal cancer-specific mortality in patients with stage ii or iii colorectal cancer: The colofol randomized clinical trial. *JAMA* 2018; 319: 2095–2103
- [115] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018; 68: 7–30
- [116] Cubiella J, Gomez R, Sanchez E et al. Endoscopic follow-up of patients after curative surgery for colorectal cancer: results of a medical assistance protocol. *Rev Esp Enferm Dig* 2003; 95: 273–281
- [117] Yun GY, Moon HS, Kwon IS et al. Left-sided colectomy: one of the important risk factors of metachronous colorectal adenoma after colectomy for colon cancer. *Dig Dis Sci* 2018; 63: 1052–1061
- [118] Matsuda T, Fujii T, Sano Y et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. *Jpn J Clin Oncol* 2009; 39: 435–442
- [119] Yoshida N, Naito Y, Siah KT et al. High incidence of metachronous advanced adenoma and cancer after endoscopic resection of colon polyps ≥ 20 mm in size. *Dig Endosc* 2016; 28: 194–202
- [120] Kim NH, Jung YS, Park JH et al. Risk of developing metachronous advanced colorectal neoplasia after colonoscopic polypectomy in patients aged 30 to 39 and 40 to 49 years. *Gastrointest Endosc* 2018; 88: 715–772
- [121] Hassan C, Quintero E, Dumonceau JM et al. Post-polypectomy colonoscopy surveillance: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2013; 45: 842–851

- 
- [122] Yazdanyar A, Newman AB. The burden of cardiovascular disease in the elderly: morbidity, mortality, and costs. *Clin Geriatr Med* 2009; 25: 563–577
- [123] Labianca R, Nordlinger B, Beretta GD et al. Early colon cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013; 24: (Suppl. 06): vi64–vi72
- [124] Watanabe T, Muro K, Ajioka Y et al. Japanese Society for Cancer of the Colon and Rectum (JSCCR) guidelines 2016 for the treatment of colorectal cancer. *Int J Clin Oncol* 2018; 23: 1–34
- [125] Benson AB 3rd, Venook AP, Al-Hawary MM et al. Rectal cancer, Version 2.2018, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Cancer Netw* 2018; 16: 874–901
- [126] Quirke P, Risio M, Lambert R et al. European guidelines for quality assurance in colorectal cancer screening and diagnosis. First Edition—Quality assurance in pathology in colorectal cancer screening and diagnosis. *Endoscopy* 2012; 44: (Suppl. 03): Se116–Se130
- [127] Poston GJ, Tait D, O’Connell S et al. Diagnosis and management of colorectal cancer: summary of NICE guidance. *BMJ* 2011; 343: d6751
- [128] Ikematsu H, Yoda Y, Matsuda T et al. Long-term outcomes after resection for submucosal invasive colorectal cancers. *Gastroenterology* 2013; 144: 551–559; quiz e514
- [129] Yoshii S, Nojima M, Noshio K et al. Factors associated with risk for colorectal cancer recurrence after endoscopic resection of T1 tumors. *Clin Gastroenterol Hepatol* 2014; 12: 292–302.e293
- [130] Backes Y, de Vos Tot Nederveen Cappel WH, van Bergeijk J et al. Risk for incomplete resection after macroscopic radical endoscopic resection of t1 colorectal cancer: a multicenter cohort study. *Am J Gastroenterol* 2017; 112: 785–796
- [131] Eckardt VF, Fuchs M, Kanzler G et al. Follow-up of patients with colonic polyps containing severe atypia and invasive carcinoma. Compliance, recurrence, and survival. *Cancer* 1988; 61: 2552–2557

CORRECTION

Hassan C, Wysocki PT, Fuccio L et al. Endoscopic surveillance after surgical or endoscopic resection for colorectal cancer: European Society of Gastrointestinal Endoscopy (ESGE) and European Society of Digestive Oncology (ESDO) Guideline

Endoscopy 2019, 51; DOI: 10.1055/a-0831-2522

In the above-mentioned article, the text at 2.3 Summary and Figure 1 have been corrected. This was corrected in the online version on February 14, 2019